

INCIDENCE OF PAPILLON-LEFÈVRE SYNDROME IN A KASHMIRI POPULATION

FAYIZAYAQOOB KHAN¹ & MUBASHIRMUSHTAQ²

¹Department of Periodontics and Oral Implantology, Government Dental College and Hospital Srinagar, Kashmir, India

²Department of Conservative Dentistry and Endodontics, Government Dental College and Hospital Srinagar, Kashmir, India

ABSTRACT

Papillon-Lefèvre syndrome is an autosomal recessive disorder with a rare occurrence. The characteristic features of this rare syndrome are palmarplantar hyperkeratosis and early onset of a severe destructive periodontitis. There is a diverse etiopathology associated with the syndrome; but a recent report has shown the syndrome is due to mutations of the cathepsin C gene. A detailed analysis was undertaken to evaluate the prevalence of this rare disorder in a Kashmiri population using clinical, laboratory and radiographic investigations.

KEYWORDS: Hyperkeratosis, Papillon-Lefèvre Syndrome & Periodontitis

Received: Apr 14, 2019; **Accepted:** May 04, 2019; **Published:** May 30, 2019; **Paper Id.:** IJDRDJUN20194

INTRODUCTION

The characteristic features of Papillon-Lefèvre syndrome (PLS) are hyperkeratosis of hands and feet and generalized aggressive periodontitis in both the primary and the permanent dentition.[1] PLS is a rare disorder with autosomal recessive inheritance. Consanguinity of parents is seen in 20% to 40% of the cases.[1] Associated clinical features include calcification of the falx cerebri and the choroid plexus.[2-4] Altered immunity to fight infections is seen in 20–25% of such patients [1,3,5,6] The recent diagnostic methods revealed the genetic defect in PLS has been mapped to chromosome 11q14–q21, which involves mutations of cathepsin C.[7,8] Studies in PLS patients display 90% reduction in cathepsin C activity.[8,9] The cathepsin C gene encodes a cysteine-lysosomal protease also known as *dipeptidyl-peptidase*. The expression of cathepsin-C is mainly seen in the epithelial regions such as palms, soles, knees, and keratinized oral gingiva. These areas are mostly affected by PLS. The etiopathology of periodontal involvement remains unknown. Periodontitis occurs after tooth eruption (10-17). There is huge plaque accumulation in the deep crevices and halitosis can occur. The primary incisors are the first teeth to show periodontitis. The whole primary dentition is shed by the age of four or five years.[1,4] Routine periodontal treatment with scaling, and root planning and other treatment modalities have proven to be of no help. [18-25].

There has not yet been any studies/case reports of this rare syndrome in Kashmiri population. The aim of this study was therefore to evaluate the frequency of Papillon-Lefèvre syndrome in a Kashmiri population using clinical, laboratory and radiological investigations.

MATERIAL AND METHODS

The study was conducted jointly in the departments of periodontics and conservative dentistry and

endodontics government dental college srinagar. A total of 10000 patients (6000 males and 4000 females) attending the general opd were included in the study. A detailed procedure was followed to evaluate each patient.

General and Extra-Oral Examination

Extra-oral examination of each patient was done to evaluate the presence of any yellowish, keratotic, confluent plaques affecting the skin of palms and soles. Examination of nails and hair were also done for each patient. Extra-oral examination of two cases revealed symmetric, well-demarcated, yellowish, keratotic, confluent plaques affecting the skin of palms and soles and extending onto the dorsal surfaces. Their nails were discolored, but hair was normal. Well circumscribed, psoriasiform, erythematous, scaly plaques were present on the knees bilaterally [Figures 1 and 2].

Intraoral examination

Intraoral examination of each case was done to evaluate any mobile and missing teeth, presence of periodontal pockets or presence of aggressive periodontitis. One case revealed missing upper left permanent central incisors, lower left permanent central and lateral incisors. Severe gingival inflammation, abscess formation, and deep periodontal pockets were noticed. Severe mobility affecting all the teeth, with heavy deposits of plaque and calculus and halitosis were also present [Figure 3]. Intraoral examination of case 2 showed deep periodontal pockets, gingival inflammation, and mobility of lower anterior teeth [Figure 4]. All primary teeth were exfoliated and no tooth was missing.

Radiographic Findings

Orthopantomographic examination of each case was done to evaluate the destruction of alveolar bone. Case 1 showed severe generalized destruction of alveolar bone. The mandibular right first molar was almost entirely out of its socket and was almost without bone support [Figure 5]. Case 2 showed generalized loss of alveolar bone [Figure 6]. On lateral skull radiograph, there was no evidence of intracranial calcification in both the patients [Figure 7].

Laboratory Investigation

Laboratory investigation was carried out, which included hematological and biochemical assessment. The results were within normal limits.



Figure 1: (a) Case 1 Presenting with Yellowish, Keratotic, Confluent Plaques Affecting the Skin of Palmar Surfaces of Hands; (b) Case 1 Presenting with Keratotic Plaques on the Dorsal Surfaces of Hands; (c) Keratotic Plaques Affecting the Dorsal Surface of Feet; (d) Several Confluent Plaques on Soles




Figure 2: (a) Case 2 Presenting with Keratotic, Confluent Plaques Affecting the Skin of Palmar Surfaces of Hands; Keratotic Plaques Affecting the Dorsal Surfaces of Hands; (c) Plaques Affecting the Dorsal Surfaces of Feet; (d) Several Confluent Plaques Affecting the Soles; (e) In Case 2, well-Circumscribed, Erythematous, Scaly Plaques on the Knees bilaterally are also Noticed



Figure 3: Intraoral Examination of Case 1 Showing Missing Upper Left Permanent

Central incisor, lower left permanent central and lateral incisors and permanent right

Lower central incisor. Severe gingival inflammation, abscess formation, and deep

Periodontal pockets were noticed. Severe mobility affecting all the teeth, with heavy

Deposits of plaque and calculus and halitosis, were also present



Figure 4: Intraoral Examination of Case 2 Showing Deep Periodontal Pockets, Gingival Inflammation, and Mobility of Lower Anterior Teeth



Figure 5: OPG of Case 1 Showing Severe Generalized Destruction of Alveolar Bone

The mandibular right first molar was almost entirely out of its socket with not much bone support



Figure 6: OPG of Case 2 Showing Generalized Destruction of Alveolar Bone

See the severe periodontal destruction in upper right first molar



Figure 7: Lateral Skull View of Case 1 Showing no Evidence of Intracranial Calcification

RESULTS

In view of the above findings, two cases were diagnosed as PLS out of 10, 000 patients with an overall percentage of 0.01% in the study population.

DISCUSSIONS

PLS has shown profound influence on the psychological and social life of the affected children. The aim of the dental clinician in such cases should be thorough dental examination and counseling of the patients parents. The etiopathology of this rare syndrome is not known some studies have proposed immunologic, microbiologic, and genetic bases. *A. actinomycetemcomitans* has been shown to be the main microorganism responsible for the aggressive periodontitis in these patients.[5,12,26,27,28] the current research has revealed the inactivation of the cathepsin C gene in

these patients.[29] In our study dermatological, periodontal, and radiological features strongly suggested the diagnosis of PLS. Consanguineous marriage of the parents of the affected children has been reported for this syndrome, [29] A definite treatment plan has not been yet devised for such patients; conventional periodontal therapy, oral hygiene instructions, and systemic antibiotics can be employed.[27] Further research is needed for evaluation and treatment of such children. Stem cell therapy can be employed in the future for the dental treatment of such children.

CONCLUSIONS

PLS threatens children and their parents with the prospect of edentulism if left untreated. Hence, early diagnosis and intervention is essential. For edentulous patients, oral rehabilitation is required; which includes partial or complete denture prosthetic replacement (according to the age of the patient). Osseointegrated implants are an option for the future and can have a great impact psychosocially by restoring esthetics as well as function. The periodontist is the first member of the health team to see and treat the patients afflicted with such unusual syndromes such as PLS, and therefore, greater awareness of this syndrome will be helpful in identifying more cases for further study.

REFERENCES

1. Papillon MM, Lefèvre P. Deux cas de keratodermie palmaire et plantaire symétrique familiale (maladie de Meleda) chez le frère et la sœur. Coexistence dans les deux cas d'altérations dentaires graves. *Bulletin de la Société Française de Dermatologie et de Syphiligraphie* 1924;31:82-7.
2. Gorlin RJ, Cohen MM, Levin LS. *Syndromes of the Head and Neck*, 3rd ed. Oxford: Oxford University Press; 1990. p. 853-5.3. Hall RK, editor. *Paediatric Orofacial Medicine and pathology*, 1st ed. London: Chapman and Hall Medical; 1994.
3. Kressin S, Herforth A, Preis S, Wahn V, Lenard HG. Papillon-Lefèvre syndrome – successful treatment with a combination of retinoid and concurrent systematic periodontal therapy: Case reports. *Quintessence Int* 1995;26:795-803.
4. Wara-Aswapati N, Lertsirivorakul J, Nagasawa T, Kawashima Y, Ishikawa I. Papillon-Lefèvre syndrome: Serum immunoglobulin G (IgG) subclass antibody response to periodontopathic bacteria. A case report. *J Periodontol* 2001;72:1747-54.
5. Lundgren T, Crossner CG, Twetman S, Ullbro C. Systemic retinoid medication and periodontal health in patients with Papillon-Lefèvre syndrome. *J Clin Periodontol* 1996;23:176-9.
6. Hart TC, Hart PS, Bowden DW, Michalec MD, Callison SA, Walker SJ, et al. Mutation of the cathepsin C gene are responsible for Papillon-Lefèvre syndrome. *J Med Genet* 1999;36:881-7.
7. Toomes C, James J, Wood AJ, Wu CL, McCormick D, Lench N, et al. Loss-of-function mutations in the cathepsin C gene result in periodontal disease and palmoplantar keratosis. *Nat Genet* 1999;23:421-4.
8. Hart PS, Zhang Y, Firatli E, Uygur C, Lotfazar M, Michalec MD, et al. Identification of cathepsin C mutations in ethnically diverse Papillon-Lefèvre syndrome patients. *J Med Genet* 2000; 37:927-32.
9. D'Angelo A, Margiotta V, Ammatuna P, Sammartano F. Treatment of prepubertal periodontitis. A case report and discussion. *J Clin Periodontol* 1992;19:214-9.
10. Brown RS, Hays G, Flaitz CM, O'Neill PA, Abramovitch K, White RR. A possible late onset variation of Papillon-Lefèvre syndrome: Report of 3 cases. *J Periodontol* 1993;64:379-86.
11. Tinanoff N, Tempore P, Maderazo EG. Dental treatment of Papillon-Lefèvre syndrome: 15 year followup. *J Clin Periodontol* 1995;22:609-12.

12. FQratlQ E, Gurel N, Efeoglu A, Badur S. Clinical andimmunological findings in 2 siblings with Papillon–Lefèvresyndrome. *J Periodontol* 1996;67:1210-5.
13. Ghaffer KA, Zahran FM, Fahmy HM, Brown RS. Papillon–Lefèvresyndrome. Neutrophil function in 15 cases from 4 familiesin Egypt. *Oral Surg Oral Med Oral Pathol Oral Radio Endod* 1999; 88:320-5.
14. Banerjee, S., & Mitra, D. (2017). *An Analytical Study on Patient Satisfaction with Respect to The Post-Traumatic Stress Disorder of Medical Practitioners*.
15. Liu R, Cao C, Meng H, Tang Z. Leukocyte function in 2 cases ofPapillon– Lefèvre syndrome. *J ClinPeriodontol* 2000; 27:69-73.
16. Lyberg T. Immunological and metabolic studies in two siblingswith Papillon– Lefèvre syndrome. *J Periodontol Res* 1982; 17:563-8.
17. Schroeder HE, Seger RA, Keller HU, Rateitschak- Plüss EM.Behavior of neutrophilic granulocytes in a case of Papillon–Lefèvre syndrome.*J ClinPeriodontol* 1983; 10:618-35.
18. Hathway R. Papillon–Lefèvre syndrome. *Br Dent J* 1982; 153:370-1.
19. Bimstein E, Lustmann J, Sela MN, Neriah ZB, Soskolne WA.Periodontitis associated with Papillon– Lefèvre syndrome.*J Periodontol* 1990;61:373-7.
20. Hattab FN, Rawashdeh MR, Yassin OM, Al-Momani AS, Al-Ubosi MM. Papillon–Lefèvre syndrome: a review of the literatureand a report of 4 cases.*JPeriodontol* 1995;66:413-20.
21. Rateitschak-Plüss EM, Schroeder HE. History of periodontitisin a child with Papillon–Lefèvre syndrome. *J Periodontol*1984;55:35 -46.
22. Glenwright HD, Rock WP. Papillon–Lefèvre syndrome.A discussion of aetiology and a case report. *Br Dent J* 1990;168:27-9.
23. Bullon P, Pascual A, Fernandez-Novoa MC, Borobio MV, Muniain MA, Camacho F. Late onset Papillon– Lefèvre syndrome?A chromosomal, neutrophil function and microbiological study.*JClinPeriodontol* 1993;20:662-7.
24. De Vree H, Steenackers K, de Boever JA. Periodontal treatmentof rapid progressive periodontitis in 2 siblings with Papillon–Lefèvre syndrome: 15- years follow-up. *J ClinPeriodontol* 2000; 27:354-60.
25. Van Dyke TE, Taubman MA, Ebersole JL, Haffajee AD,Socransky SS, Smith DJ, et al. The Papillon–Lefèvre syndrome:Neutrophil dysfunction with severe periodontal disease. *Clin Immunol Immunopathol* 1984;31:419-29.
26. Saidunnisa, B., Atiqulla, S., & Ayman, G. Prevalence of Polycystic Ovarian Syndrome among Students of RAK Medical and Health Sciences University United Arab Emirates.
27. Sagllie FR, Marfany A, Camargo P. Intragingival occurrenceof Actinobacillusactinomycetemcomitans and Bacterioidesgingivalis in active destructive periodontal lesions. *J Periodontol*1988;59:259-65.
28. Stabholz A, Taichman S, Soskolne WA. Occurrence ofactinobacillusactinomycetemcomitans and anti-leukotoxinantibodies in some members of an extended family affected byPapilon–Lefèvre syndrome. *J Periodontol* 1995; 66:653-7.
29. Firatli E, Gurel N, Efeoglu A, Badur S. Clinical and immunologicalfindings in 2 siblings with Papillon-Lefèvre syndrome.*J Periodontol* 1996;67:1210-5.
30. Hart TC, Hart PS, Bowden DW, Michalec MD, Callison SA,Walker SJ, et al. Mutations of the cathepsin C gene are responsiblefor Papillon-Lefèvre syndrome. *J Dent Res* 2004; 83:368-70.